

1

Which of the following is the area of the triangle whose vertices are the points

which represent the cubic roots of unity on Argand's plane?

- $\frac{3\sqrt{3}}{4}$

- $\frac{\sqrt{3}}{2}$

- $\frac{3\sqrt{3}}{2}$

- $\frac{\sqrt{3}}{4}$

# 2

The number of ways can a person in a sport club participates in 3 games

at least from the set { football , hand ball , volley ball , basket ball } equals .....

- ${}^4C_3 + {}^4C_4$

- ${}^4C_3 \times {}^4C_4$

- ${}^4P_3 + {}^4P_4$

- ${}^4P_3 \times {}^4P_4$

# 3

In the expansion of  $\left(x^2 - \frac{1}{x}\right)^{15}$  according to the descending powers of  $x$ ,  
the value of the term free of  $x$  equals .....

- $^{15}C_5$
- $-^{15}C_5$
- $^{15}C_9$
- $-^{15}C_9$

# 4

If the two planes:

$18x + 15y - 6z + 1 = 0$ ,  $ax + by + 2z + 1 = 0$  are parallel,

then  $ab = \dots\dots\dots$

- 30
- -30
- 90
- -90

# 5

If  $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$  then  $(A^2)^{-1} = \dots\dots\dots$

•  $\begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix}$

•  $\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$

•  $\begin{pmatrix} -\cos \theta & -\sin \theta \\ -\sin \theta & -\cos \theta \end{pmatrix}$

•  $\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix}$

# 6

If the straight line  $\frac{x - 2}{3} = \frac{y + 1}{-4} = \frac{z + 3}{5}$  makes with the planes  $x y$ ,  $y z$ ,  $z x$  angles of measures

L, M, N respectively, then  $\sin^2 L + \sin^2 M + \sin^2 N = \dots\dots\dots$

- 1
- 2
- $\sqrt{3}$
- $\frac{3}{2}$

# 7

If  $1, \omega, \omega^2$  are the cubic roots of unity, then  $\left(\frac{a}{\omega} - \frac{a}{\omega^2} + \frac{3a}{\omega^4} - \frac{3a}{\omega^5}\right)^2 = \dots\dots\dots$

- $-48 a^2$
- $48 a^2$
- $16 a^2$
- $-16 a^2$

# 8

In the expansion of  $\left( a x^2 - \frac{b}{x} \right)^{12}$  according to the descending powers of  $x$ .

$T_7$  is .....

- The term containing  $x^6$
- The term free of  $x$
- The term before the last
- The term containing  $x^7$



# 9

The code of a lock consists of 3 different digits number chosen from the digits { 1 , 2 , 3 , ..... , 9 } .

By how many ways we can form a code contains the digit 6 ?

- 168
- 126
- 336
- 224

# 10

If the coefficient of the ninth term in the expansion of  $\left(a\sqrt{x} - \frac{1}{a\sqrt{x}}\right)^{12}$

according to the descending power of X equals 7920 , then a = .....

- $\pm \frac{1}{2}$
- $\pm 2$
- $\pm \frac{1}{4}$
- $\pm 4$

# 11

$$\begin{vmatrix} x & y & y \\ y & x & y \\ y & y & x \end{vmatrix} = (x + 2y) \times \dots\dots\dots$$

$$\bullet \begin{vmatrix} 1 & y & y \\ 0 & x-y & 0 \\ 0 & 0 & x-y \end{vmatrix}$$

$$\bullet \begin{vmatrix} 1 & y & y \\ 0 & x+y & 0 \\ 0 & 0 & x+y \end{vmatrix}$$

$$\bullet \begin{vmatrix} 1 & y & 0 \\ 0 & x+y & 0 \\ 0 & 0 & x-y \end{vmatrix}$$

$$\bullet \begin{vmatrix} 1 & y & y \\ 0 & x-y & 2y \\ 0 & 0 & x+y \end{vmatrix}$$

# 12

If  $A(3, -4, 0)$ ,  $B(15, 0, 2)$ ,  $C(0, -8, 4)$  are three points in the space and they are the vertices of  $\triangle ABC$ , then the distance between the centroid point of the  $\triangle ABC$  and the plane  $XZ$  is .....

- greater than its distance from the plane  $xy$
- greater than its distance from the plane  $yz$
- smaller than or equal to its distance from the plane  $xy$
- greater than or equal to its distance from the plane  $yz$

# 13

The possible values of K which make the distance between the two points

A ( 2 , K , 3 ) , B ( -4 , 4 , 2 ) equals  $\sqrt{62}$  are .....

- -1 or 9
- -5 or -9
- 1 or 5
- 1 or -9

# 14

If the shortest distance between the point A ( 3 , 5 , 1 ) and the surface of the sphere whose centre M ( 1 , 2 , - 5 ) is 2 length unit , then the radius of the sphere = ..... length unit.

- 5
- 2
- 7
- 12

# 15

If the measure of the angle between the two planes :

$\vec{r} \cdot (3, -4, 2) = 7$  and  $3x + 4y - mz = 12$  is  $90^\circ$ , then  $m = \dots\dots\dots$

•  $\frac{-7}{2}$

•  $\frac{-3}{2}$

•  $\frac{-25}{2}$

•  $\frac{3}{2}$

# 16

If  $Z_1 = 3 (\cos 300^\circ + i \sin 300^\circ)$ ,  $Z_2 = 2 (\sin 240^\circ + i \cos 240^\circ)$

, then which of the following represents the exponential form of  $Z_1 Z_2$ ?

- $6 e^{\frac{5}{6} \pi i}$
- $6 e^{\pi i}$
- $\frac{3}{2} e^{\frac{5}{6} \pi i}$
- $\frac{3}{2} e^{\pi i}$



# 17

If  $2 \cdot {}^{n+1}C_r = {}^{n+1}P_r$ ,  $\frac{{}^nC_{r+1}}{{}^nC_r} = \frac{5}{3}$

, then  ${}^nC_r + {}^nP_r = \dots\dots\dots$

- 63
- 33
- 60
- 36

# 18

If  $\vec{A}$ ,  $\vec{B}$  are two vectors where  $\|\vec{A}\| = 5$ ,

and the component of vector  $\vec{B}$  in the direction of vector  $\vec{A}$  is 3, then  $\vec{A} \cdot \vec{B} = \dots\dots\dots$

• 15

•  $\frac{5}{3}$

•  $\frac{3}{5}$

• 8

# 19

If  $a e^{2\theta i} + b e^{-2\theta i} = 5 \cos 2\theta - i \sin 2\theta$  where  $a, b$  are two positive real numbers,

$\theta \in ]0, \frac{\pi}{2}[$ ,  $i^2 = -1$ , then:  $a b = \dots\dots\dots$

- 6
- 2
- 5
- 3

# 20

If  ${}^{n+1}P_r > {}^{n+1}P_{r-1}$ , then  $n > \dots\dots\dots$

- $r - 1$
- $r - 3$
- $r + 1$
- $1 - r$

# 21

If the greatest coefficient in the expansion of  $(a + x)^{20}$  is the coefficient of  $T_{11}$ ,

then  $a \in \dots\dots\dots$  where  $a \in R^+$

•  $\left[ \frac{10}{11}, \frac{11}{10} \right]$

•  $[10, 11]$

•  $\left[ \frac{-11}{10}, \frac{10}{11} \right]$

•  $\left[ \frac{9}{11}, \frac{10}{11} \right]$

If  $A^*$  is the augmented matrix for the linear system of equations

$3x + 2y - z = 4$ ,  $x + y - z = 3$ ,  $x = 2z$ , then .....

- $2 < \text{Rk}(A^*) < 4$
- $\text{Rk}(A^*) < 3$
- $1 < \text{Rk}(A^*) \leq 2$
- $1 \leq \text{Rk}(A^*) < 3$

# 23

If  $\vec{A}$ ,  $\vec{B}$ ,  $\vec{C}$  represent three adjacent edges in a parallelepiped,  $\|\vec{A}\| = 2$

and the direction angles of vector  $\vec{A}$  are  $(135^\circ, 60^\circ, 120^\circ)$ ,  $\vec{B} = (1, \sqrt{2}, 0)$ ,  $\vec{C} = (\sqrt{2}, 3, 5)$ ,

then the volume of the parallelepiped = ..... cubic unit .

- 16
- $6\sqrt{2}$
- 11
- $16\sqrt{2}$

If the plane  $2x - y + 2z = 6$  touches the surface of the sphere whose equation  $x^2 + y^2 + z^2 - 4x - 2y + 6z + 5 = 0$ , then the equation of the straight line which passing through the center of the sphere and the point of tangency is .....

- $\vec{r} = (2, 1, -3) + t(2, -1, 2)$
- $\vec{r} = (2, 1, -3) + t(4, 0, -1)$
- $\vec{r} = (4, 0, -1) + t(2, 1, -3)$
- $\vec{r} = (2, -1, 2) + t(2, 1, 3)$



If the plane  $bcx + acy + abz = abc$  intersects the coordinate axes  $x$ ,  $y$  and  $z$  at the points  $K$ ,  $N$  and  $M$  respectively and the plane  $bcx + acy - abz = -abc$  intersects the coordinate axes  $x$ ,  $y$  and  $z$  at the points  $K'$ ,  $N'$  and  $M$  respectively, then the pyramid  $MKNK'/N'$  is .....

(where  $a$ ,  $b$ ,  $c$  are positive real numbers and  $a \neq b$ )

- right quadrilateral pyramid.
- regular quadrilateral pyramid.
- right triangular pyramid.
- regular triangular pyramid.